

FORM PTO-1449
(Modified)U.S. Department of Commerce
Patent and Trademark Office

Attorney Docket No.: UM-06106

Serial No.: 09/778,496

INFORMATION DISCLOSURE STATEMENT BY APPLICANT
(Use Several Sheets if Necessary)Applicant: David M. Lubman *et al.*

(37 CFR § 1.98(b))

Filing Date: 02/07/01

Group Art Unit:

U.S. PATENT DOCUMENTS

Examiner Initials	Cite No.	Serial / Patent Number	Issue Date	Applicant / Patentee	Class	Subclass	Filing Date
CSM	1	5498545	March 12, 1996	Vestal	436	47	July 21, 1994
↓	2	5670054	September 23, 1997	Kibbey <i>et al.</i> ,	210	656	April 4, 1996

FOREIGN PATENTS OR PUBLISHED FOREIGN PATENT APPLICATIONS

		Document Number	Publication Date	Country / Patent Office	Class	Subclass	Translation	
							Yes	No
CSM	3	EP0617048	1994-09-28	PCT	—	—		
↓	4	WO9701755	1997-01-16	PCT	—	—		
	5	WO0158925	2001-08-16	PCT	—	—		
	6	WO0158926	2001-08-16	PCT	—	—		
↓	7	WO 98/40395	1998-09-17	PCT	—	—		
	8	WO02088701	2002-07-11	PCT	—	—		

OTHER DOCUMENTS (Including Author, Title, Date, Relevant Pages, Place of Publication)

CSM	9	Andrews, <i>et al.</i> , Analysis of DNA adducts using high-performance separation techniques coupled to electrospray ionization mass spectrometry, <i>J Chromatogr A</i> , 856(1-2):515 (1999);						
↓	10	Ball and Mascagni, Purification of synthetic peptides using reversible chromatographic probes based on the Fmoc molecule, <i>Int J Pept Protein Res</i> , 40(5):370 (1992);						
	11	Davidsson and Nilsson, Peptide mapping of proteins in cerebrospinal fluid utilizing a rapid preparative two-dimensional electrophoretic procedure and matrix-assisted laser desorption/ionization mass spectrometry, <i>Biochim Biophys Acta</i> , 1473(2-3):391 (1999);						
	12	Lee, Protein separation using non-porous sorbents, <i>J Chromatogr B Biomed Sci Appl</i> , 699(1-2):29 (1997);						
	13	Medzihradszky <i>et al.</i> , Protein sequence and structural studies employing ionization-high energy collision-induced dissociation, <i>International J of Mass Spectrometry and Ion Processes</i> , 160(1):357 (1997);						
	14	Nimura <i>et al.</i> , Fast Protein separation by reversed-phase high-performance liquid chromatography on octadecylsilyl-bonded non-porous silica gel effect of particle efficiency, <i>Journal of Liquid Chromatography</i> , 585(2):207 (1991);						
↓	15	Richmond <i>et al.</i> , High-throughput flow injection analysis-mass spectrometry with networked delivery of colour rendered results: the characterisation of liquid chromatography fractions, <i>Journal of Chromatography</i> , 835(1-2):29 (1999);						
	16	Hermann and Andreas, Mapping and identification of <i>Corynebacterium glutamicum</i> proteins by two-dimensional gel electrophoresis and microsequencing, <i>Electrophoresis</i>, 19(18):3217 (1998);						
	17	Houen and Bach, Characterization of protein carboxy-terminal ends using carboxypeptidase peptide Y: Sequence, composition, and identification of the carboxy terminal peptide by peptide mapping, <i>Methods Molecular Cellular Biology</i>, 3(4):175 (1992);						
CSM	18	Patterson and Aebersold, Mass spectrometric approaches for the identification of gel-separated proteins, <i>Electrophoresis</i> , 16(10):1791 (1995);						
	19	Raznikov <i>et al.</i>, Selective digital filtering of mass spectra of chromatography data for determination of "target" compounds in complex mixtures, <i>Advances in Mass Spectrometry</i>, 14(E0442804): (1998);						
	20	SZE and Dominic, Time of flight effects in matrix assisted laser desorption/ionization Fourier transform mass spectrometry, <i>Rapid Commun Mass Spectrom</i>, 13(5):390 (1999);						
CSM	21	Wall, <i>et al.</i> , Isoelectric focusing nonporous silica reversed-phase high-performance liquid chromatography/electrospray ionization time-of-flight mass spectrometry: a three-dimensional liquid-phase protein separation method as applied to the human erythroleukemia cell-line, <i>Rapid Commun Mass Spectrom</i> , 15(18):1649 (2001);						
↓	22	Wall, <i>et al.</i> , Three-dimensional protein map according to pI, hydrophobicity and molecular mass, <i>J Chromatogr B Analyt Technol Biomed Life Sci</i> , 774(1):53 (2002);						

Examiner:

Date Considered:

EXAMINER:

Initial citation considered. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

FORM PTO-1449 (Modified)		U.S. Department of Commerce Patent and Trademark Office		Attorney Docket No.: UM-06106	Serial No.: 09/778,496
INFORMATION DISCLOSURE STATEMENT BY APPLICANT (Use Several Sheets If Necessary)				Applicant: David M. Lubman <i>et al.</i>	
				Filing Date: 02/07/01	Group Art Unit:
(37 CFR § 1.98(b))					
OTHER DOCUMENTS (Including Author, Title, Date, Relevant Pages, Place of Publication)					
	23	Zgoda, <i>et al.</i> , Prosovsky, Prediction and experimental confirmation of the cytochrome b5 three-dimensional peptide map, Physical Chemical Biology and Medicine, 2(3):135 (1995);			
CJM	24	Bini, <i>et al.</i> , Protein expression profiles in human breast ductal carcinoma and histologically normal tissue, Electrophoresis, 18(15):2832 (1997);			
	25	Chen, <i>et al.</i> , Identification of proteins from two-dimensional gel electrophoresis of human erythroleukemia cells using capillary high performance liquid chromatography/electrospray-ion trap-reflectron time-of-flight mass spectrometry with two-dimensional topographic map analysis of in-gel tryptic digest products, Rapid Commun Mass Spectrom, 13(19):1907 (1999);			
	26	Clauser, <i>et al.</i> , Role of accurate mass measurement (+/- 10 ppm) in protein identification strategies employing MS or MS/MS and database searching, Anal Chem, 71(14):2871 (1999);			
	27	Dai, <i>et al.</i> , Two-layer sample preparation: a method for MALDI-MS analysis of complex peptide and protein mixtures, Anal Chem, 71(5):1087 (1999);			
	28	Damerval, Quantification of silver-stained proteins resolved by two-dimensional electrophoresis: genetic variability as related to abundance and solubility in two maize lines, Electrophoresis, 15(12):1573 (1994);			
✓	29	Fuqua, <i>et al.</i> , Induction of the estrogen-regulated "24K" protein by heat shock, Cancer Res, 49(15):4126 (1989);			
	30	Hanash, Advances in Electrophoresis, 1:44 (1998);			
CJM	31	Herbert, Advances in protein solubilisation for two-dimensional electrophoresis, Electrophoresis, 20(4-5):660 (1999);			
	32	Hoogland C., The 1999 SWISS-2DPAGE database update., Nucleic Acids Res, 28:286 (2000);			
	33	Immmer, <i>et al.</i> , Identification of phosphorylated proteins from thrombin-activated human platelets isolated by two-dimensional gel electrophoresis by electrospray ionization-tandem mass spectrometry (ESI-MS/MS) and liquid chromatography-electrospray ionization-mass spectrometry (LC-ESI-MS), Electrophoresis, 19(6):1015 (1998);			
	34	Kahn, From genome to proteome: looking at a cell's proteins, Science, 270(5235):369 (1995);			
	35	Matsui, <i>et al.</i> , Immobilized pH gradient two-dimensional gel electrophoresis and mass spectrometric identification of cytokine-regulated proteins in ME-180 cervical carcinoma cells, Electrophoresis, 18(3-4):409 (1997);			
	36	Mohammad, <i>et al.</i> , Induced expression of alpha-enolase in differentiated diffuse large cell lymphoma, Enzyme Protein, 48(1):37 (1994);			
	37	Neidhardt, <i>et al.</i> , Genomically linked cellular protein databases derived from two-dimensional polyacrylamide gel electrophoresis, Electrophoresis, 10(2):116 (1989);			
	38	Neubauer and Mann, Mapping of phosphorylation sites of gel-isolated proteins by nanoelectrospray tandem mass spectrometry: potentials and limitations, Anal Chem, 71(1):235 (1999);			
	39	O'Farrell, High resolution two-dimensional electrophoresis of proteins, J Biol Chem, 250(10):4007 (1975);			
	40	Opitck, <i>et al.</i> , Comprehensive two-dimensional high-performance liquid chromatography for the isolation of overexpressed proteins and proteome mapping, Anal Biochem, 258(2):349 (1998);			
	41	Patterson, Matrix-assisted laser-desorption/ionization mass spectrometric approaches for the identification of gel-separated proteins in the 5-50 pmol range, Electrophoresis, 16(7):1104 (1995);			
	42	Rasmussen, <i>et al.</i> , Two-dimensional gel database of human breast carcinoma cell expressed proteins: an update, Electrophoresis, 19(5):818 (1998);			
	43	Redner, <i>et al.</i> , The t(5;17) variant of acute promyelocytic leukemia expresses a nucleophosmin-retinoic acid receptor fusion, Blood, 87(3):882 (1996);			
	44	Reid, <i>et al.</i> , Capillary column chromatography improves sample preparation for mass spectrometric analysis: complete characterization of human alpha-enolase from two-dimensional gels following in situ proteolytic digestion, Electrophoresis, 19(6):946 (1998);			
✓	45	Reymond, <i>et al.</i> , Standardized characterization of gene expression in human colorectal epithelium by two-dimensional electrophoresis, Electrophoresis, 18(15):2842 (1997);			
	46	Righetti, Isoelectric Focusing: Theory, Methodology, and Applications, in Laboratory Techniques in Biochemistry and Molecular Biology, v. 11, Ed. Burdon R.H., Elsevier Biomedical Press, (1983);			
CJM	47	Rosenfeld, <i>et al.</i> , In-gel digestion of proteins for internal sequence analysis after one- or two-dimensional gel electrophoresis, Anal Biochem, 203(1):173 (1992);			
Examiner:		Gey Mah		Date Considered: September 25, 2003	
EXAMINER: Initial citation considered. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant					

U.S. Department of Commerce
Patent and Trademark Office

Serial No.: 09/778,496

Applicant: David M. Lubman *et al.*

Filing Date: 02/07/01

Group Art Unit:

OTHER DOCUMENTS (Including Author, Title, Date, Relevant Pages, Place of Publication)

[illegible]

C. M. 1 f

September 25, 2003

EXAMINER:

Initial citation considered. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.